

POSTER PRESENTATION

Open Access

Detection of infection with *Angiostrongylus vasorum* (Nematoda, Strongylida) by PCR

Mohammad Al-Sabi^{1*}, Pia Webster², Jacob Willesen³, Peter Deplazes⁴, Alexander Mathis⁴, Christian Kapel¹

From Parasite infections of domestic animals in the Nordic countries – emerging threats and challenges. The 22nd Symposium of the Nordic Committee for Veterinary Scientific Cooperation (NKVet) Helsinki, Finland. 7-9 September 2008

Background

The French heart worm *Angiostrongylus vasorum* is a parasitic nematode of the pulmonary arteries and heart of canines often with severe and in some cases fatal outcome. The diagnosis is based on detection and species identification of larvae in faeces which can be problematic in Veterinary praxis especially in cases with low excreting animals. A reliable technique is thus needed for correct diagnosis and estimation of the true prevalence of infection in a population as well as for monitoring and control campaigns.

Materials and methods

A PCR was developed from the ITS2 region of the rDNA of *A. vasorum*. The sensitivity of the primers was tested with DNA from adult *A. vasorum* from a naturally infected fox and first stage larvae (L1) from an experimentally infected foxes. The specificity of the primers was tested against DNA from the most common helminth parasites of canines in Denmark and neighbouring countries. Furthermore the PCR system was applied as a confirmative test in a screening study of Danish hunting dogs and an epidemiological study of helminth parasites of wildlife in Denmark.

Results

The designed primers were very sensitive and could detect a single *A. vasorum* L1. The primers were also very specific and did not react with DNA from any of the common canine helminths. When used as a confirmative test, the PCR system proved to be robust and easy to work with detecting a single larva, and for use in post mortem examination of wildlife. There are practical

problems that can face the PCR system such as isolating dead larvae from frozen samples and the known problem of intermittent larval excretion in dogs. These two problems can be solved by isolation of larvae by sieving instead of by Baermann sedimentation if samples were frozen, and examining consecutive fresh faecal samples.

Conclusions

We were able to design a new PCR to detect DNA of *A. vasorum* in canines. The test proved to be very sensitive and specific when tested in clinical and epidemiological studies. The test will be further applied in many epidemiological and clinical studies to come.

Author details

¹Department of Agriculture and Ecology, University of Copenhagen, DK-1871 Frederiksberg C, Denmark. ²Department of Disease Biology, University of Copenhagen, DK-1871 Frederiksberg C, Denmark. ³Small animal hospital, University of Copenhagen, DK-1871 Frederiksberg C, Denmark. ⁴Institute of Parasitology, University of Zurich, CH-8057 Winterthurerstrasse 266A, Zurich, Switzerland.

Published: 13 October 2010

doi:10.1186/1751-0147-52-S1-S9

Cite this article as: Al-Sabi et al.: Detection of infection with *Angiostrongylus vasorum* (Nematoda, Strongylida) by PCR. *Acta Veterinaria Scandinavica* 2010 **52**(Suppl 1):S9.

¹Department of Agriculture and Ecology, University of Copenhagen, DK-1871 Frederiksberg C, Denmark

Full list of author information is available at the end of the article