

Experimental Infection of Conventional Pigs with *Streptococcus suis* serotype 2 by Aerosolic Exposure

By L. W. Madsen¹, B. Nielsen², B. Aalbæk³, H. E. Jensen¹, J. P. Nielsen⁴ and H. J. Riising²

¹Department of Pharmacology and Pathobiology, ³Department of Veterinary Microbiology, and ⁴Department of Clinical Studies, Royal Veterinary and Agricultural University, Copenhagen, and ²Intervet Scandinavia AS, Skovlunde, Denmark.

Streptococcus suis is an important cause of meningitis, arthritis and septicaemia, especially in young pigs. Serotype 2 is the most prevalent type in clinical material from pigs in Europe (Wisselink *et al.* 2000) and the infection causes severe disease outbreaks in swine herds. Different experimental models have been used to elucidate the infection but central parts of the pathogenesis still remain unclear (Gottschalk & Segura 2000). In spontaneous infection, *S. suis* is generally believed to invade via the upper respiratory tract (Gottschalk & Segura 2000). Recently, we described an infection model in minipigs using aerogenous challenge and subsequent steroid treatment (Madsen *et al.* 2001a). In conventional pigs, reports of models attempting aerosol exposure seem limited to a single study using anaesthetized pigs (Chengappa *et al.* 1986).

In order to evaluate the pathogenesis of *S. suis* type 2 infection in pigs, we aimed at establishing an aerosol model for the infection using unanaesthetized conventionally reared, weaned pigs. The present report describes the microbiological and pathological findings in a pilot study of an aerosol model for *S. suis* infection in conventional pigs.

Four clinically healthy, 6-week-old, female,

Landrace-Yorkshire crossbred pigs (animals A-D) from the same litter (weaned at app. 4 weeks of age) were included in this study. They were obtained from a herd with no history of disease compatible with *S. suis* infection and *S. suis* had never been isolated from the herd. For the aerosol exposure, a 1.5 m³ chamber connected to a nebulizing apparatus was used. The chamber and procedure used were essentially as previously described (Madsen *et al.* 2001a). Briefly, the animals were in the chamber for 20 min while 40 ml of an aquatic solution of 1% acetic acid (pH 3.4) was dispersed and let into the chamber with atmospheric air. Then the pigs were kept outside the chamber for one hour. Thereafter, 3 of the animals (A-C) were brought back into the chamber and 42.5 ml of a bacterial suspension was dispersed over 20 min in the air supply to the chamber. The bacterial suspension was a 5-h broth culture of *S. suis* serotype 2 (strain P321/6) concentrated to 1.1×10^{10} colony-forming units/ml. The fourth pig (D) served as a control to evaluate the immediate effect of acetic acid alone. One h after exposure to acetic acid in the chamber, this animal was euthanized by exsanguination after being anaesthetized by intramuscular injection (1 ml/15 kg) of Zoletil® (25 mg/ml zolezepam,

25 mg/ml tiletamin). The 3 remaining animals were housed in a single pen and observed for clinical signs of disease and rectal temperatures were recorded daily. On the fifth day after exposure, animal A was euthanized due to the severity of clinical signs. The 2 remaining pigs were euthanized on the sixth day.

Following euthanasia, all pigs were necropsied and gross lesions were recorded. Tissues for microscopy as well as swabs for microbiological culturing were collected to cover all parts of the respiratory tract and a range of organs known to be affected in *S. suis* infection (Madsen et al. 2001a). From each animal, samples were taken from gross lesions as well as a standard of 31 tissues for histopathology and 18 tissues for microbiological culture, which was done aerobically at 37°C on 5% calf blood agar. Morphologically suspect colonies were subcultured and identified biochemically and serolog-

ically using standard methods. From aseptically sampled blood as well as from areas with a normal bacterial flora, i.e., nasal cavity, pharyngeal and palatine tonsils, microbiological culture was done as previously described (Madsen et al. 2001a).

For histopathology, fixation in 4% neutral buffered formaldehyde, decalcification of relevant tissues, as well as processing of HE stained sections were performed as previously described (Madsen et al. 1998). The presence of *S. suis* antigen was examined by immunohistochemistry using a previously published protocol (Madsen et al. 2001b).

Clinical signs were only observed in pigs A and C with onset on days 3 and 5 after exposure, respectively. These pigs had loss of appetite, elevated body temperatures (40.0-41.5°C), were reluctant to rise and lame in one or more legs. Gross lesions were noted in animal A, in which

Table 1. Distribution of lesions and of *S. suis* serotype 2 as detected by culture and immunohistochemistry in 3 animals after aerosolic challenge.

Tissue	Animals positive		Characteristic macroscopic and/or histologic lesions (Animals)
	Culture	IHC	
Nasal cavity	C	A,C	Mild, focal mucopurulent rhinitis (A,C)
Lung	-	A	Diffuse fibrinopurulent pleuritis (A,B); acute embolic pneumonia (A)
Bronchial lnn.	-	A	Follicular hyperplasia and exudation of heterophils (A,B)
Middle ear	C	B	Unilateral purulent otitis media and exudate in Eustachian tube (B)
Inner ear	n.d.	C	Bilateral purulent labyrinthitis and perineuritis (n. vestibularis) (C) - Fig. 1
Tonsil, pharyngeal	A,C	C	Focal exudation of heterophils (C)
Tonsil, soft palate	A,B,C	A,B,C	Heterophils in crypt epithelium and lumen (A,B,C)
Brain	C	C	Diffuse fibrinopurulent meningitis with focal submeningeal encephalitis (C)
Heart	A	A	Diffuse fibrinopurulent pericarditis (A)
Peritoneum	A	A	Diffuse fibrinopurulent peritonitis (A)
Mesenteric lnn.	-	A	Follicular hyperplasia and exudation of heterophils (A)
Liver	A	A	Multifocal, subcapsular hepatic necrosis and fibrinopurulent perihepatitis (A)
Spleen	C	A	-
Joints	A,C	A,C	Suppurative arthritis (A,C)
Blood	A	n.d.	-

n.d.: not done; -: negative; IHC: immunohistochemistry for *S. suis* serotype 2; A,B,C: animal no.



Figure 1. Inner ear, *S. suis* infected pig (animal C). Suppurative labyrinthitis with exudate (arrow) in the perilymph of the cochlear scala tympani. HE $\times 4$.

a fibrinopurulent polyserositis was seen, and in animal C, which had an exudative meningitis and arthritis.

By microscopy, a range of lesions of a generally suppurative nature were noted in the challenged animals (Table 1). Lesions compatible with septicaemia were seen in animal A. In animal C, the histopathological findings included suppurative meningitis, arthritis, and labyrinthitis (Fig. 1). In the third *S. suis* exposed animal, B, a suppurative otitis media was observed histologically. In the control animal, D, the only changes observed were focal, mild mucopurulent rhinitis and focal, mild, chronic, purulent bronchopneumonia in the right cranial lung lobe.

By immunohistochemistry, *S. suis* serotype 2 antigen was demonstrated in animals A-C in a number of tissues (Table 1). Generally, antigen detection was related to the presence of 1-3 μm

structures, resembling coccoid bacteria, located intracellularly or in exudates. Rarely, a more diffuse intracellular immunostaining was observed in phagocytes in inflamed tissues or corresponding lymphoid tissues. *S. suis* serotype 2 was reisolated from various tissues in animals A, B, and C (Table 1). No bacterial pathogens were isolated from animal D.

The clinical signs as well as the lesions observed in 2 of the 3 animals challenged with *S. suis* were comparable to spontaneous *S. suis* serotype 2 infection in pigs (Reams *et al.* 1994). Acute embolic pneumonia was also observed in animal A. Although this observation is not common in *S. suis* infections, bacterial embolism is not surprising given the bacteraemia present. In animal C, a bilateral labyrinthitis with *S. suis* antigen present in the exudate was detected. This lesion, which apparently has not been seen previously in experimental pig models of this infection, is a common sequela in porcine meningitis due to spontaneous *S. suis* serotype 2 infection (Madsen *et al.* 2001b). Furthermore, hearing loss due to inner ear affection is a major characteristic in humans affected by *S. suis* meningitis (Arends & Zanen 1988). In animal B, otitis media with *S. suis* antigen present was observed. As no signs of a systemic infection were observed in this animal, these findings might constitute a local infection after direct spread via the likewise affected Eustachian tube.

In the control animal D, only minor lesions were found. Thus, the observed focal bronchopneumonia is regarded as being insignificant. However, the mucopurulent rhinitis also seen in the infected animals may have been associated with the exposure to acetic acid. This would be in concordance with previous studies, where intranasal instillation of 1% acetic acid was shown to induce loss of cilia and epithelium as well as submucosal oedema and inflammation within 12 h of exposure (Gagné & Mar-

tineau-Doizé 1993). However, further studies are required to evaluate the effect of exposure to acetic acid and the interaction with *S. suis* infection in this model.

In conclusion, by experimental aerosolic exposure, infection with *S. suis* serotype 2 was established in unanaesthetized conventional pigs, and lesions similar to those seen in spontaneously infected animals were induced.

Acknowledgments

We wish to thank Dr. T. Gonzales for her collaboration and appreciate the technical assistance from A. Wamsler-Jensen, L. Johansen, P.R. Mortensen, L. Kioerboe, E. Hjuler and M. Bak. This study was financially supported by the Danish Bacon and Meat Council and by a R N Thomsen grant.

References

- Arends JP, Zanen HC: Meningitis caused by *Streptococcus suis* in humans, *Rev. Infect. Dis.*, 1988, 10, 131-137.
- Chengappa MM, Maddux RL, Kadel WL, Greer SC, Herren CE: *Streptococcus suis* infection in pigs: Incidence and experimental reproduction of the syndrome. Proc. 29th ann. meeting, Amer. Assn. Vet. Lab. Diagnosticians, Louisville, Kentucky, 1986, pp. 25-38.
- Gagné S, Martineau-Doizé B: Nasal epithelial changes induced in piglets by acetic acid and by *Bordetella bronchiseptica*, *J. Comp. Pathol.*, 1993, 109, 71-81.
- Gottschalk M, Segura M: The pathogenesis of the meningitis caused by *Streptococcus suis*: the unresolved questions, *Vet. Microbiol.*, 2000, 76, 259-272.
- Madsen LW, Aalbaek B, Nielsen OL, Jensen HE: Aerogenous infection of microbiologically defined minipigs with *Streptococcus suis* serotype 2 – a new model, *APMIS*, 2001a, 109, in press.
- Madsen LW, Jensen AL, Larsen S: Spontaneous lesions in clinically healthy, microbiologically defined Göttingen minipigs, *Scand. J. Lab. Anim. Sci.*, 1998, 25, 159-166.
- Madsen LW, Svensmark B, Elvestad K, Jensen HE: Otitis interna is a frequent sequela to *Streptococcus suis* meningitis in pigs, *Vet. Pathol.*, 2001b, 38, 190-195.
- Reams RY, Glickman LT, Harrington DD, Thacker HL, Bowersock TL: *Streptococcus suis* infection in swine: a retrospective study of 256 cases. Part II. Clinical signs, gross and microscopic lesions, and coexisting microorganisms, *J. Vet. Diagn. Invest.*, 1994, 6, 326-334.
- Wisselink HJ, Smith HE, Stockhofe-Zurwieden N, Peperkamp K, Vecht U: Distribution of capsular types and production of muramidase-released protein (MRP) and extracellular factor (EF) of *Streptococcus suis* strains isolated from diseased pigs in seven European countries, *Vet. Microbiol.*, 2000, 74, 237-248.

(Received January 22, 2001; accepted February 15, 2001).

Reprints may be obtained from: L.W. Madsen, Laboratory of Veterinary Pathology, Department of Pharmacology and Pathobiology, Royal Veterinary and Agricultural University, Copenhagen, 17 Bülowsvej, DK-1870 Frederiksberg C, Denmark. E-mail: lwm@kvl.dk, tel:+45 35 28 31 16, fax: +45 35 28 31 11.